Proceedings of the Second Annual LHCP

August 26, 2014

Measurement of hadron composition in charged jets from pp collisions with the ALICE experiment

arXiv:1408.5723v1 [hep-ex] 25 Aug 2014

Benjamin Andreas Hess [1]

On behalf of the ALICE Collaboration,
Physikalisches Institut, University of Tübingen, Germany

ABSTRACT

We report on the first measurement of the charged hadron composition in charged jets from pp collisions. The ALICE detector at the LHC was used to study charged pion, kaon and (anti-)proton production in jets. The results were extracted from 2×10^8 minimum bias events at a centre-of-mass energy of $\sqrt{s} = 7\,\mathrm{TeV}$.
We present the transverse momentum (p_T) spectra and reduced momentum spectra ($z^\mathrm{ch} \equiv p_\mathrm{T}^\mathrm{track}/p_\mathrm{T,jet}^\mathrm{ch}$) of π, K and p in charged jets with $p_\mathrm{T,jet}^\mathrm{ch}$ between 5 and $20\,\mathrm{GeV}/c$. The measurements are compared to Monte Carlo calculations.

PRESENTED AT

The Second Annual Conference
on Large Hadron Collider Physics
Columbia University, New York, U.S.A
June 2-7, 2014

[1] *Email address*: Benjamin-Andreas.Hess@Uni-Tuebingen.de.
Work supported by BMBF project 05P12VTCAA.

1 Introduction

Jets are phenomenological objects constructed to represent partons originating from hard scattering processes. The present knowledge about parton fragmentation into identified hadrons is mainly constrained by jet fragmentation measurements at e^+e^- colliders [1, 2]. In addition, neutral jet fragments have been measured by the CDF Collaboration [3]. The ALICE experiment [4] at the LHC has powerful particle identification (PID) capabilities allowing for the measurement of identified charged hadron spectra in jets from pp collisions for the first time. The charged hadron composition is extracted with sophisticated PID techniques using the specific energy loss ($\mathrm{d}E/\mathrm{d}x$) of tracks in the Time Projection Chamber (TPC) [5]. The TPC is the main tracking and PID device of ALICE and has a $\mathrm{d}E/\mathrm{d}x$ resolution of about 5% for pp collisions. The measurement presented here is based on the ALICE analysis strategy for the measurement of inclusive charged particle production in charged jets [6], but with particle identification using the TPC.

2 Analysis technique

The analysis is carried out on a sample of 2×10^8 minimum bias (MB) events from pp collisions at a centre-of-mass energy of $\sqrt{s} = 7\,\mathrm{TeV}$ recorded with the ALICE detector in 2010. It is based on charged tracks with transverse momentum $p_\mathrm{T} > 0.15\,\mathrm{GeV}/c$ and within the pseudo-rapidity range $|\eta^\mathrm{track}| < 0.9$, which are reconstructed with the ALICE Inner Tracking System (ITS) and the TPC. The tracks are clustered to jets using the anti-k_T algorithm of the FastJet [7] package with resolution parameter $R = 0.4$. Only jets with $|\eta^\mathrm{jet}| < 0.5$, that are fully contained in the ALICE central barrel acceptance, are included in the analysis.

Based on the TPC $\mathrm{d}E/\mathrm{d}x$, the raw π, K and p yields differential in $p_\mathrm{T}^\mathrm{track}$ or $z^\mathrm{ch} \equiv p_\mathrm{T}^\mathrm{track}/p_\mathrm{T,jet}^\mathrm{ch}$ are extracted. This extraction is done separately for $p_\mathrm{T,jet}^\mathrm{ch} = 5-10$, $10-15$ and $15-20\,\mathrm{GeV}/c$.

Finally, the raw yields are corrected for detector efficiency, acceptance, (jet) p_T resolution, secondary particle and muon contamination. The latter correction is required because the TPC $\mathrm{d}E/\mathrm{d}x$ resolution does not allow for the separation of pions and muons, i.e. the muon yield is attributed to the pion yield. The correction procedure is based on that for the inclusive charged particle measurement described in [6] and has been extended to take into account particle type dependent effects. The following discussion will focus on the PID in jets.

The raw π, K and p differential yields are extracted with the TPC Coherent Fit [8]. The TPC Coherent Fit is a 2-dimensional fitting procedure that analyses the TPC $\mathrm{d}E/\mathrm{d}x$ distribution as a function of particle momentum (p). It is able to extract raw particle yields differentially in p_T from $0.15\,\mathrm{GeV}/c$ to above $20\,\mathrm{GeV}/c$ with an accuracy of better than 10% on average for the ALICE pp collision data. The TPC Coherent Fit is based on the observation that the mean and width of the $\mathrm{d}E/\mathrm{d}x$ signal as well as the particle fractions of each species are continuous as a function of particle momentum. Using continuous $\mathrm{d}E/\mathrm{d}x$ models and allowing only for statistical fluctuations of the particle fractions in neighbouring bins, it simultaneously extracts the particle yields and the $\mathrm{d}E/\mathrm{d}x$ model parameters in a single optimisation procedure.

Technically, the TPC Coherent Fit maximises a log-likelihood function l,

$$l \equiv \tilde{l} + l^\mathrm{reg}, \tag{1}$$

where the additional term l^reg is a regularisation term for the particle yields and will be discussed below. If f_{ki} denotes the particle fraction of species k in momentum bin i and $s_k(p_i;\vec{\Theta})$ is the $\mathrm{d}E/\mathrm{d}x$ model for species k at momentum p_i with *a priori* unknown parameters $\vec{\Theta}$, the log-likelihood term $\tilde{l}$ symbolically reads

$$\tilde{l} \equiv \sum_i l_\mathrm{i} \left(\sum_k f_{ki} s_k(p_i;\vec{\Theta}) \right), \tag{2}$$

where l_i is the log-likelihood function for momentum bin i*. The sum $\sum_i$ adds up the contributions from different momentum bins, whereas the sum $\sum_k$ represents the superposition of the $\mathrm{d}E/\mathrm{d}x$ distributions

*The parameter is the modelled $\mathrm{d}E/\mathrm{d}x$ distribution in that momentum bin. The likelihood that the model describes the measured distribution is evaluated with Poissonian statistics.

from different particle species. The functional form of the particle fraction f_{ki} is not known, but a continuity condition on the particle momentum can be imposed that only allows for statistical deviations of the particle fraction from the interpolated value from the neighbouring momentum bins. This is achieved by adding the regularisation term

$$l^{\mathrm{reg}} \equiv \sum_{k,i} l^{\mathrm{reg}}_{ki}(f_{ki}) \tag{3}$$

to $\tilde{l}$ in Eq. 1, where the regularisation strength contributes equally for each particle species and momentum bin[†].

With the $\mathrm{d}E/\mathrm{d}x$ distribution as a function of momentum as input, the TPC Coherent Fit maximises the log-likelihood in Eq. 1 yielding both the $\mathrm{d}E/\mathrm{d}x$ model parameters $\vec{\Theta}$ and the particle fractions f_{ki}. The single optimisation procedure is driven *coherently* by the full-range constraint of the $\mathrm{d}E/\mathrm{d}x$ models and the constraint on the particle fractions.

Utilising the $\mathrm{d}E/\mathrm{d}x$ for PID at high particle momentum ($p > 4\,\mathrm{GeV}/c$) is challenging because the extracted particle yields are highly sensitive to the mean $\mathrm{d}E/\mathrm{d}x$. Since the K-p separation at such momenta is about 5% in ALICE pp data, a 1‰ mean $\mathrm{d}E/\mathrm{d}x$ bias is estimated to cause a 2% bias of the particle fractions. Hence, extensive studies of the systematic uncertainties of the TPC Coherent Fit have been performed. These include $\mathrm{d}E/\mathrm{d}x$ model uncertainties, robustness against changes of the $\mathrm{d}E/\mathrm{d}x$ quality, a possible particle type dependence of the $\mathrm{d}E/\mathrm{d}x$ (which is found to be negligible) and $p^{\mathrm{ch}}_{\mathrm{T,jet}}$ dependence of the $\mathrm{d}E/\mathrm{d}x$. The latter is caused by an increased local track density in jets and automatically taken into account by the fit, since the $\mathrm{d}E/\mathrm{d}x$ model and fractions are fitted in each $p^{\mathrm{ch}}_{\mathrm{T,jet}}$ bin separately.

In addition to these studies, the TPC Coherent Fit was applied to Monte Carlo (MC) samples (PYTHIA [9] tune Perugia0 [10]) with full ALICE detector simulation (with GEANT3 [11]) and reconstruction in the same way as done for data. Within uncertainties (typically smaller than 10%), the results of this procedure reproduce the MC truth at the detector level.

Furthermore, the results of the TPC Coherent Fit have been cross-checked by an independent method—the TPC Multi-Template Fit. The TPC Multi-Template Fit extracts in great detail the TPC $\mathrm{d}E/\mathrm{d}x$ response up to intermediate momenta ($p \lesssim 8\,\mathrm{GeV}/c$) from pure MB data samples selected via TPC $\mathrm{d}E/\mathrm{d}x$, Time-Of-Flight (TOF) and track topology (products of $\mathrm{K}^0_{\mathrm{S}}$, Λ and $\overline{\Lambda}$ decays and of γ conversions). The $\mathrm{d}E/\mathrm{d}x$ response at high momenta is determined from model fits to these clean samples. With this response, templates for the $\mathrm{d}E/\mathrm{d}x$ distributions of each species are generated for the considered input data sample. Finally, the particle fractions are estimated by minimising the difference between the measured $\mathrm{d}E/\mathrm{d}x$ distribution and the sum of templates weighted with the particle fractions. The template generation and fitting is performed directly in p_{T} or z^{ch} bins, the only free parameters being the particle fractions in each bin. As for the TPC Coherent Fit, the TPC Multi-Template fit uses a log-likelihood maximisation with a regularisation term for the particle fractions.

The two methods have different sources of systematic uncertainties. As shown in Fig. 1, the results of both methods agree within uncertainties. It can also be seen that PYTHIA Perugia0 MC deviates from data beyond the systematic error.

3 Results

The fully corrected p_{T} differential yields per jet of π, K and p in charged jets are shown in Fig. 2. The spectra span $3-4$ orders of magnitude and become harder with increasing $p^{\mathrm{ch}}_{\mathrm{T,jet}}$. A clear $p^{\mathrm{ch}}_{\mathrm{T,jet}}$ ordering of the spectra is observed; the ordering inverts at $p_{\mathrm{T}} = 0.4\,\mathrm{GeV}/c$ for π and at $2\,\mathrm{GeV}/c$ for K and p.

In Fig. 3, the fully corrected K/π and p/π ratios in charged jets are shown as a function of z^{ch}. The K/π ratio exhibits a monotonic increase with z^{ch}, reaching $0.5-0.6$, indicating that the strangeness fraction in jets rises with z^{ch}. For the p/π ratio, a maximum of $0.15-0.2$ is reached at $z^{\mathrm{ch}} = 0.5-0.6$ followed by a decrease as z^{ch} approaches unity. This indicates that leading baryons in charged jets are suppressed.

[†]The regularisation term l^{reg}_{ki} is derived from a Gaussian likelihood and, thus, has a proper statistical interpretation.



Figure 1: Uncorrected K/π ratio as a function of z^{ch} for $p_{\mathrm{T,jet}}^{\mathrm{ch}} = 10 - 15\,\mathrm{GeV}/c$ for data and MC (PYTHIA Perugia0). The uncorrected results for data of the TPC Coherent Fit (full points) are compared to those of the TPC Multi-Template Fit (rectangles) and the detector level MC truth (open points). The error bars represent the statistical uncertainties and the error boxes indicate the systematic uncertainties.



Figure 2: Corrected p_{T} spectra of π (*left*), K (*middle*) and p (*right*) in charged jets from pp collisions at $\sqrt{s} = 7\,\mathrm{TeV}$. The spectra for $p_{\mathrm{T,jet}}^{\mathrm{ch}} = 5 - 10$ (diamonds), $10 - 15$ (circles) and $15 - 20\,\mathrm{GeV}/c$ (stars) are shown.

Comparing the ratios for the three different $p_{\mathrm{T,jet}}^{\mathrm{ch}}$ bins, a scaling is observed for $z^{\mathrm{ch}} > 0.2$ in all $p_{\mathrm{T,jet}}^{\mathrm{ch}}$ bins for the K/π ratio and for $p_{\mathrm{T,jet}}^{\mathrm{ch}} = 10 - 15$ and $15 - 20\,\mathrm{GeV}/c$ in case of p/π.

In Fig. 4, the spectra shown in Fig. 2 are compared to the PYTHIA [9] tunes Perugia0, Perugia0NoCR (*noCR* stands for *no colour reconnection*) and Perugia2011 [10] for π, K and p. The best agreement is observed at high $p_{\mathrm{T,jet}}^{\mathrm{ch}}$ and high particle p_{T}. For low particle p_{T}, all considered PYTHIA tunes undershoot (overshoot) the pions (protons). Typically, all three tunes describe the data within 30% except for protons with $p_{\mathrm{T}} < 0.5\,\mathrm{GeV}/c$, where the deviation goes beyond 100%, and with p_{T} close to the upper bound of the $p_{\mathrm{T,jet}}^{\mathrm{ch}}$ bin, where the discrepancy is around 50% for some tunes. The maximum of the proton spectra around $p_{\mathrm{T}} = 2\,\mathrm{GeV}/c$ (cf. Fig. 2) is reproduced very well by all PYTHIA tunes, but they fail to describe the width and the high-p_{T} slope. Of the tunes considered, Perugia0NoCR gives the best description of the K spectra in all $p_{\mathrm{T,jet}}^{\mathrm{ch}}$ bins.



Figure 3: Corrected K/π (*left*) and p/π (*right*) ratios as a function of z^{ch} in charged jets from pp collisions at $\sqrt{s} = 7\,\mathrm{TeV}$. The ratios for $p_{\mathrm{T,jet}}^{\mathrm{ch}} = 5 - 10$ (diamonds), $10 - 15$ (circles) and $15 - 20\,\mathrm{GeV}/c$ (stars) are shown.



Figure 4: MC/data ratios of the π (*left*), K (*middle*) and p (*right*) p_{T} spectra in charged jets from pp collisions at $\sqrt{s} = 7\,\mathrm{TeV}$.

4 Conclusions

We presented the first measurement of identified jet fragmentation of charged hadrons at hadron colliders from ALICE. The particle yields and ratios as functions of p_{T} and z^{ch} of primary hadrons (π, K, p) in charged jets from pp collisions at $\sqrt{s} = 7\,\mathrm{TeV}$ with $p_{\mathrm{T,jet}}^{\mathrm{ch}} = 5 - 20\,\mathrm{GeV}/c$ are extracted using advanced PID techniques. We observe that the $p_{\mathrm{T,jet}}^{\mathrm{ch}}$ scaling of the z^{ch} spectra disappears at lowest $p_{\mathrm{T,jet}}^{\mathrm{ch}}$. Furthermore, our measurements show an increase of the strangeness fraction with z^{ch} and a suppression of leading baryons at high z^{ch}. PYTHIA simulations reproduce the data typically within 30% accuracy. However, we observe a tension between data and PYTHIA at low $p_{\mathrm{T,jet}}^{\mathrm{ch}}$ and for pions and protons at low particle p_{T}.

References

[1] J. P. Lees *et al.* [BaBar Collaboration], Phys. Rev. D **88** (2013) 3, 032011.

[2] M. Leitgab *et al.* [Belle Collaboration], Phys. Rev. Lett. **111** (2013) 6, 062002.

[3] T. Aaltonen *et al.* [CDF Collaboration], Phys. Rev. D **88** (2013) 9, 092005.

[4] K. Aamodt *et al.* [ALICE Collaboration], JINST **3** (2008) S08002.

[5] J. Alme, Y. Andres *et al.*, Nucl. Instrum. Meth. A **622** (2010) 78 316.

[6] O. Busch for the ALICE Collaboration, arXiv:1306.2747 [nucl-ex].

[7] M. Cacciari, G. P. Salam and G. Soyez, Eur. Phys. J. C **72** (2012) 1896.

[8] X.-G. Lu, Doctoral Dissertation at the University of Heidelberg (2013), CERN-THESIS-2013-179.

[9] T. Sjostrand, S. Mrenna and P. Z. Skands, JHEP **0605** (2006) 026.

[10] P. Z. Skands, Phys. Rev. D **82** (2010) 074018.

[11] R. Brun, F. Carminati and S. Giani, "GEANT – Detector Description and Simulation Tool", CERN-W5013, 1993.